Wisconsin AIDS/HIV Surveillance Summary Cases reported 1983 through 2006

BROWN COUNTY

| | | ulative <u>Percent</u> | - | I 2001-2005 <u>Percent</u> | | d in 2006 <u>Percent</u> | Presum <u>Cases</u> | ned alive Percent |
|--|---------------|---------------------------|----|-------------------------------|----|-----------------------------|------------------------|----------------------|
| Total cases | 313 | 100.0% | 64 | 100.0% | 12 | 100.0% | 190 | 100.0% |
| Current disease category | | | | | | | | |
| AIDS | 218 | 69.6% | 38 | 59.4% | 6 | 50.0% | 108 | 56.8% |
| Non-AIDS | 95 | 30.4% | 26 | 40.6% | 6 | 50.0% | 82 | 43.2% |
| Risk Exposure Categories | | | | | | | | |
| Men who have sex with men (MSM) | 190 | 60.7% | 37 | 57.8% | 11 | 91.7% | 114 | 60.0% |
| Injecting drug use (IDU) | 30 | 9.6% | 6 | 9.4% | 0 | 0.0% | 21 | 11.1% |
| MSM and IDU | 30 | 9.6% | 5 | 7.8% | 1 | 8.3% | 15 | 7.9% |
| Hemophilia/Coagulation disorder | 8 | 2.6% | 0 | 0.0% | 0 | 0.0% | 1 | 0.5% |
| High-risk heterosexual contact | 24 | 7.7% | 8 | 12.5% | 0 | 0.0% | 17 | 8.9% |
| Transfusion-associated | 5 | 1.6% | 0 | 0.0% | 0 | 0.0% | 2 | 1.1% |
| Mother with/at risk | 3 | 1.0% | 1 | 1.6% | 0 | 0.0% | 3 | 1.6% |
| Undetermined | 23 | 7.3% | 7 | 10.9% | 0 | 0.0% | 17 | 8.9% |
| Gender | | | | | | | | |
| Female | 39 | 12.5% | 12 | 18.8% | 0 | 0.0% | 32 | 16.8% |
| Male | 274 | 87.5% | 52 | 81.3% | 12 | 100.0% | 158 | 83.2% |
| Unknown | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| Race/Ethnicity | | | | | | | | |
| White | 257 | 82.1% | 43 | 67.2% | 11 | 91.7% | 147 | 77.4% |
| Black | 23 | 7.3% | 9 | 14.1% | 0 | 0.0% | 18 | 9.5% |
| Hispanic | 21 | 6.7% | 11 | 17.2% | 0 | 0.0% | 15 | 7.9% |
| Asian/Pacific Islander | 2 | 0.6% | 0 | 0.0% | 0 | 0.0% | 2 | 1.1% |
| American Indian | 10 | 3.2% | 1 | 1.6% | 1 | 8.3% | 8 | 4.2% |
| Multi-racial | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| Unknown | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| Age at diagnosis* | | | | | | | | |
| Under 5 | 3 | 1.0% | 1 | 1.6% | 0 | 0.0% | 0 | 0.0% |
| 5-14 | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 2 | 1.1% |
| 15-19 | 6 | 1.9% | 1 | 1.6% | 0 | 0.0% | 0 | 0.0% |
| 20-24 | 28 | 8.9% | 7 | 10.9% | 1 | 8.3% | 1 | 0.5% |
| 25-44 | 226 | 72.2% | 46 | 71.9% | 10 | 83.3% | 101 | 53.2% |
| 45 years and older | 50 | 16.0% | 9 | 14.1% | 1 | 8.3% | 85 | 44.7% |
| Unknown | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| * For cases presumed alive, age is the age | on 12/31/2006 | | | | | | | |
| Year of Report | | | | | | | | |
| Before 1990 | 45 | 14.4% | | | | | | |
| 1990 | 12 | 3.8% | | | | | | |
| 1991 | 19 | 6.1% | | | | | | |
| 1992 | 20 | 6.4% | | | | | | |
| 1993 | 20 | 6.4% | | | | | | |
| 1994 | 18 | 5.8% | | | | | | |
| 1995 | 31 | 9.9% | | _ | | | | |
| 1996 | 16 | 5.1% | | | | | | |
| 1997 | 23 | 7.3% | | | | | | |
| 1998 | 12 | 3.8% | | | | | | |
| 1999 | 6 | 1.9% | | | | | | |
| 2000 | 15 | 4.8% | | | | | | 7 |
| 2001 | 13 | 4.2% | | | | | | , |
| 2002 | 17 | 5.4% | | | | | | ١ |
| 2003 | 10 | 3.2% | | | | | | |
| 2004 | 11 | 3.5% | | | | | | - |
| 2005 | 13 | 4.2% | | | | | | |
| 2006 | 12 | 3.8% | | | | | | |